



ELSEVIER

REVIEW ARTICLE

The Epithelial Gatekeeper Against Food Allergy

Linda Chia-Hui Yu

Graduate Institute of Physiology, National Taiwan University College of Medicine, Taipei, Taiwan

Received: May 14, 2009

Revised: Jul 27, 2009

Accepted: Jul 29, 2009

KEY WORDS:

allergen uptake;

commensal bacteria;

epithelial permeability;

food allergy;

IgE/CD23;

intestinal barrier function

The rapid rise of allergic disorders in developed countries has been attributed to the hygiene hypothesis, implicating that increased environmental sanitation in early childhood may be associated with higher incidence of hypersensitivity. Intestinal epithelial barriers play a crucial role in the maintenance of gut homeostasis by limiting penetration of luminal bacteria and dietary allergens, yet allowing antigen sampling via the follicle-associated epithelium for generation of tolerance. However, this intricate balance is upset in allergic intestines, whereby luminal proteins with antigenic properties gain access to the subepithelial compartment and stimulate mast cell degranulation. Recent studies demonstrated that food allergens were protected from lysosomal degradation, and were transported in large quantities across the epithelium by binding to cell surface IgE/CD23 (FcεRII) that prevented the antigenic protein from lysosomal degradation in enterocytes. IL-4 (a Th2-type cytokine) not only increased production of IgE from B cells, but also upregulated the expression of CD23 on intestinal epithelial cells. Further studies indicated that CD23 was responsible for the bidirectional transport of IgE across epithelium. The presence of IgE/CD23 opens a gate for intact dietary allergens to transcytose across the epithelial cells, and thus foments the mast cell-dependent anaphylactic responses. The understanding of the molecular mechanism responsible for epithelial barrier defects may be helpful in designing novel therapies to treat food allergy and other allergic diseases.

1. Clinical Prevalence of Food Allergies

Increasing prevalence of allergic diseases, such as asthma, hay fever, eczema and food allergies, has been observed in recent decades. The number of visits due to food-induced anaphylaxis is growing in hospital emergency departments and has significantly increased the burden on medical systems.^{1,2} Food allergy is reported in 6–10% of the pediatric population, and is more frequent in children than adults.^{3,4} The most common allergens include cow's milk, eggs, peanuts, and seafood. Cow's milk allergy occurs in 2–3% of the infant population, and is mostly diagnosed within the first month of life. Egg allergy is seen in 1–2% and peanut allergy in 0.3% of children.

Although some types of food allergy, e.g. cow's milk, remit spontaneously during the first few years of life, they are often associated with later development of respiratory and skin allergic manifestations.^{3,4}

2. Hygiene Hypothesis

The rapid rise of allergic disorders in developed countries has led to the postulation that lower incidence of microbial infection in early childhood—due to increased environmental sanitation—may be associated with higher incidence of hypersensitivity.⁵ This is termed the hygiene hypothesis. This hypothesis was originally formed in 1989 by Strachan,

Corresponding author. Graduate Institute of Physiology, National Taiwan University College of Medicine, 1 Jen-Ai Road, Section 1, Taipei 100, Taiwan.
E-mail: lchyu@ntu.edu.tw

an epidemiologist, who observed an inverse correlation between family size and the occurrence of atopic diseases.⁶ Speculations were formed that smaller household size indicated less cross-infection between family members/siblings—this could be related to the development of allergic disorders in individuals. This phenomenon was coined the “post-industrial revolution epidemic”. In the following years, allergists and immunologists further broadened the concept that reduced exposure to microorganisms and microbial products, and increased inoculation of antibiotics and vaccines at a young age may predispose individuals to allergic diseases.^{6,7}

3. Oral Tolerance and Allergy

Oral tolerance has been defined as a systemic immune unresponsiveness to a specific antigen which was previously orally administered. Studies in experimental models show that the dose and frequency of antigen feeding influences the course of tolerance acquisition.^{8,9} The putative concept is that feeding a high dosage of antigen results in clonal deletion or anergy of specific T cell clones that involve Fas/FasL-dependent apoptosis, whereas feeding a low dosage of antigen favors the pathway of active suppression following the induction of regulatory T (Treg) cells in the Peyer's patches (PP). However, it should be kept in mind that the two forms of tolerance are not mutually exclusive but are likely to be overlapped in physiological situations.

PPs are specialized lymphoid follicles in the gut that are implicated as a crucial site for induction of oral tolerance. Luminal antigens are readily taken up by follicle-associated epithelium (mainly by M cells) on PP and delivered to the underlying dendritic cells and lymphocytes for trafficking to mesenteric lymph nodes (MLN).^{10,11} Previous studies demonstrated that lymphotoxin (LT) α knock-out mice, or mice treated *in utero* with LT β receptor-blockade, are devoid of PP and MLN, and fail to develop oral tolerance.^{12,13} However, mesenteric lymph nodes have been suggested to be responsible for the induction of high-dose oral tolerance in the absence of PP. An elegant study showed that by treating pregnant LT α knock-out mice with an agonist anti-LT β R monoclonal antibody, their progeny developed MLN but without PP and readily acquired systemic tolerance to the orally administered protein.^{14,15}

Following antigen feeding, the presence of antigen-specific CD4+CD25+Treg cells and secretion of suppressive cytokines, e.g. transforming growth factor (TGF) β , are found in the PP compartment.^{16–19} The involvement of TGF β in oral tolerance induction was first suggested in studies using oral feeding

of myelin basic protein to suppress experimental autoimmune encephalomyelitis.²⁰ Neutralizing antibody to TGF β inhibited the bystander suppression of immune responses conferred by tolerized T cells *in vitro*.²⁰ Further evidence indicated that TGF β played a complementary role in facilitating the induction of low dose antigen-induced oral tolerance by CD4+CD25+Treg cells and was crucial for the maintenance of established tolerance.^{21,22} Moreover, oral administration of TGF β along with antigen augments the reduction of antigen-specific immunoglobulin levels, T-cell reactivity, and immediate skin hypersensitivity compared with antigen feeding alone.²²

Treg cells mediate active suppression and are primary effector cells involved in the generation of oral tolerance. Oral administration of ovalbumin to ovalbumin TCR transgenic mice resulted in a relative increase in the population of CD4+CD25+Tregs expressing CTLA-1. Adoptive transfer of these CD4+CD25+Tregs from antigen-fed ovalbumin TCR transgenic mice suppressed the delayed type hypersensitivity responses in recipient mice, suggesting that this particular cell type is responsible for oral tolerance induction to specific antigens.²³ Recent data indicated that a dendritic cell subset in the lamina propria characterized by the expression of CD103 induced naive T cells to differentiate into CD4+CD25+Foxp3+Treg cells, via a TGF β and retinoic acid-dependent mechanism.^{24,25}

Dysregulation of various functions in PP may promote allergic sensitization. Studies using milk allergic mice have shown that enhanced transfer of aggregates through PP or abnormal interaction between lymphocytes and dendritic cells in PP favors the development of food allergy.^{18,26,27} Orally administered aggregated proteins were redirected to uptake through PP which promoted a significant Th2 response and triggered allergic sensitization in a cow's milk-allergic mouse model.²⁷ Interestingly, subsequent anaphylaxis was only elicited by orally given soluble, but not aggregated, forms of allergen. The findings suggested that the transport of soluble protein via villous epithelial cells was the main pathway for anaphylactic responses.²⁷

In healthy individuals, the colonic lumen hosts over 100 trillion commensal bacteria forming the enteric microfloral population, and is established after birth.²⁸ Enteric bacteria are involved in numerous protective and metabolic functions of the gut, such as preventing colonization of pathogens, degrading non-digestible dietary substances, producing short chain fatty acids and vitamins, and shaping the mucosal immunity.^{28–31} The intestinal IgA production is profoundly affected by the colonization of commensal microflora, as shown by the low levels of IgA in germ-free animals, which is corrected after

inoculation of luminal bacteria.³² Luminal antigen sampling by PP induces class switch recombination of IgM-positive B cells to IgA. The classing switching of IgA is partially dependent on presence of TGF β and production of retinoic acid by intestinal dendritic cells.^{33,34} These IgA-committed B cells drain to the mesenteric lymph nodes, subsequently enter the thoracic duct and bloodstream, and finally home back to the intestinal mucosa. The secretory IgA (sIgA) produced by these lamina propria plasma cells is transported across the intestinal epithelium via the polymeric immunoglobulin receptor into the lumen.³⁵ Enteric bacteria are coated with sIgA.³⁶ Recent studies showed that luminal sIgA selectively adhered to M cells in the mouse and human intestinal PP via a novel IgA receptor—this also mediated translocation of bacteria and antigenic products to the underlying dendritic cells.^{37,38} The bacteria transported by the sIgA across M cells further induces naïve B cells to differentiate into IgA-committed plasma cells.³⁹ The roundtrip, bidirectional transport of sIgA and sIgA-mediated bacterial coatings have been implicated in the mechanism of antigen neutralization that curtails bacterial overgrowth.^{40,41}

Recent evidence indicated that gut bacterial flora is required for the induction of oral tolerance. In contrast to conventionally-raised animals, germ-free mice do not generate immune tolerance against fed antigen.^{42,43} Mice given oral antibiotics during infancy showed an increase in serum levels of IgG1 and IgE, and decreased IgG2a levels that are associated with enhanced IL-4 secretion in stimulated spleen cells, suggesting a Th2-polarized immune response.⁴⁴ The absence of T cells in PP under germ-free conditions was implicated in the mechanism of failure of tolerance induction.⁴⁵

4. Immunological Basis of Food Allergy

Food allergy is characterized by sensitization to specific dietary antigens and local inflammation at the encountering site of allergens. The onset of sensitization involves the combination of a number of factors including genetic traits, allergen exposure, intestinal microflora, and environmental stimuli. Recent evidence has suggested that psychological stress also predisposes animals to sensitization against orally given antigens which otherwise induced tolerance.⁴⁶

The sensitization phase of allergy is characterized by the production of IgE and Th2-type cytokine (IL-4, IL-5, and IL-13) responses. Elevated production of IL-4 by mononuclear cells has been demonstrated in the blood and intestinal mucosa of atopic individuals.^{47,48} IL-4 induces germ line ϵ transcript for isotype class switching in B cells and

promotes B cell proliferation to increase the synthesis of antigen-specific IgE.⁴⁹ In addition to its presence in serum, elevated levels of IgE are detected in the intestinal fluid in food-allergic patients, suggesting that IgE is also secreted into the gut lumen.^{50,51} In parasite-infected rats with stimulated IgE production, the concentration of IgE in the intestinal fluid was greater than that in the serum or mesenteric lymph.⁵² A previous study demonstrated that radiolabeled IgE (but not IgG) was translocated from the serum into the gut wall and lumen after infusion of rats with IL-4.⁵³

IgE binding to the high affinity Fc ϵ RI on the surface of mast cells represents the hallmark of allergy. Cross-linking of IgE by specific antigen induces mast cell degranulation and release of mediators, thereby causing anaphylactic responses.⁵⁴ Anaphylactic reactions in food allergy are associated with enhanced epithelial ionic transport with passive out flux of water—this is responsible for clinical diarrheal symptoms.^{55,56} Mast cell mediators, e.g. histamine and prostaglandin, are involved in the stimulation of epithelial ion secretion.^{57,58} Histamine evokes chloride secretion in epithelial cell monolayers through H1-receptors on intestinal epithelial cells, causing elevation in cytosolic calcium level and protein kinase C activation.^{59,60} Prostaglandins have been shown to increase the secretory response of intestinal epithelial monolayers directly via cAMP upregulation, or to serve as a mediation step for ion secretion induced by several cytokines and mediators, such as TNF α , serotonin, leukotrienes, and proteases.^{61–63}

5. Epithelial Barrier Defects in Intestinal Allergy

It is generally accepted that intestinal anaphylactic reactions are caused by biological mediators released from mucosal mast cells after antigen cross-linking of IgE bound to the cell surface. However, macromolecular food antigens must first cross the intestinal epithelial barrier before gaining access to mast cells in the subepithelial compartment. Abnormal antigen passage through PP has been suggested as one of the mechanisms responsible for the lack of tolerance.^{18,26,27} In comparison to the limited exposing region of PPs, villous epithelium constitutes a relatively large surface area and plays a more important role in responsible for the breach of barrier in intestinal allergy.

Physiologically, most dietary proteins are digested by luminal gastric and pancreatic proteases as well as integral brush border enzymes. They are converted to small peptides and amino acids, which are then absorbed by enterocytes via electrogenic or

sodium-dependent transporters. Although a small amount of intact protein is endocytosed by intestinal epithelial cells in normal conditions, most of it is sorted into lysosomal compartments for degradation and therefore, transcytosis of whole proteins are prevented.^{64–66} In other words, the transcellular route across epithelial cells contributes partially to exclusion of luminal proteins with antigenic properties. The intestinal epithelial cells are joined at their apical side by tight junctions.^{67,68} The transmembranous junctional proteins, e.g. claudins, occludin or junction-associated molecules, are linked to intracellular zonula occludens which are bridges to cytoskeletal actin and myosin filaments. The tight junctional complexes are the rate-limiting step for paracellular diffusion, and only permit small molecules lesser than 500 daltons to cross between cells. Focal disorganization of tight junctional proteins or circumferential contraction of the perijunctional actomyosin ring may increase paracellular permeability.^{69–73}

The place in which route food allergens cross the gut epithelium has been extensively explored in the last decade. In rodent models, it was observed that addition of antigen from the luminal or serosal side of the allergic intestine induces strong ion secretory responses, with different timeframes. Antigen challenge to the serosal side of the intestine induces an immediate increase (approximately 30 seconds) in ion secretion, whereas luminal addition of antigen

results in a lag phase (approximately 3 minutes) before the occurrence of the secretory response, mediated by mast cell activation.⁵⁷ The lag phase of the anaphylactic response after luminal challenge appears to reflect the time for antigen transport across the intestinal epithelial cells to activate mast cells in the lamina propria.⁵⁷

In contrast to protein degradation in normal intestinal epithelial cells, enhanced rates of transcytosis of intact proteins was documented in allergic animals.^{74,75} Previous studies modeling luminal allergen challenge demonstrated that increased antigen uptake was observed within the endosomal compartment of jejunal enterocytes in sensitized rats before mast cell activation.^{74,76,77} Studies using sensitized Ws/Ws rats (mast cell-deficiency due to mutation of the c-kit gene) and mast cell stabilizing agents provided evidence that mast cell activation does not play a role in the mechanism of heightened apical-to-basolateral transcellular transport of antigen at this timepoint.⁷⁶ The uptake of antigen appeared to be specific and the transport pathway was exclusively transcellular within the first 2 minutes post-challenge. This period of specific transcellular antigen transport before mast cell activation was termed phase I.^{76,77} The period following mast cell activation by epithelial ion secretory response was termed phase II (Figure 1). During phase II, antigens were visualized not only inside endosomes but also within the tight junctions and paracellular




Figure 1 Intestinal epithelial barrier defects in food allergy. Physiologically, most dietary proteins are digested to small peptides and amino acids before being absorbed into enterocytes. A very small percentage of intact proteins may be endocytosed into epithelial cells but are degraded by lysozymes and lose their antigenic properties. In food allergy, IL-4 (a Th2-type cytokine) induces B cells to produce large amounts of IgE to be secreted into serum and gut lumen, or surface-bound onto mast cells. IL-4 also upregulates the expression of low affinity IgE receptor, CD23 (FcεRII), on intestinal epithelial cells. Following exposure to dietary allergens, enhanced transepithelial antigen transport is mediated by IgE/CD23 prior to mast cell activation during phase I. Transcytosed allergens reach the subepithelial lamina propria and cause IgE cross-linking on mast cells, resulting in cell degranulation and anaphylactic responses. The release of mast cell mediators, such as histamine, prostaglandin and proteases, are known to induce epithelial ion secretion and increased paracellular epithelial permeability in phase II.

regions between enterocytes in allergic rats.^{76,78} The electrical conductance (measurement of ionic permeability through the paracellular pathway) in intestinal tissues of allergic rats was comparable to those of non-sensitized control animals during phase I. This suggests that gut paracellular permeability was not modified in response to sensitization *per se*. Moreover, a gradual time-dependent increase in tissue conductance corresponds to the phenomenon of enhanced paracellular antigen transport in allergic rats during phase II. The abnormal paracellular epithelial permeability in phase II was absent in allergic mast cell-deficient Ws/Ws rats, suggesting a crucial role of mast cell activation in the induction of tight junction opening and increase of paracellular influx that was not antigen-specific.

Allergic animal models have shown that enhanced transepithelial antigen transport prior to mast cell activation is specific for the allergen to which the rodents are sensitized, suggesting an immunoglobulin recognition mechanism at the level of the epithelium.^{74,75} A low affinity IgE receptor, CD23/Fc ϵ RII, was previously known for its role in regulating IgE synthesis in B cells and promoting B cell proliferation.^{79–81} The expression of CD23 was also found in small intestinal epithelial cells in normal and food allergic humans and rodents, as well as in bronchial epithelial cells in asthmatic patients.^{75,77,82} Studies in sensitized rat intestines have demonstrated the translocations of CD23 from cell surface to the membrane of allergen-containing endosomes, confirming the internalization of CD23 protein upon luminal antigen challenge.⁷⁵ Further studies using genetically mutant mouse models provided evidence for the role of IgE/CD23 in mediating enhanced transepithelial antigen transport in allergy.^{77,78} The phenomenon of augmented antigen uptake in allergic enterocytes was completely absent in sensitized CD23^{−/−} mice and IL-4^{−/−} mice. Moreover, the increased transepithelial antigen uptake in allergic wild type mice was inhibited luminally with neutralizing anti-CD23 antibodies.^{77,78} Passive sensitization of naive mice by injecting immune serum from allergic mice restored the allergic response, but not if IgE was first depleted from serum, confirming the crucial role of IgE in antigen uptake. In addition, IL-4 increased the expression of CD23 transcript and protein levels in murine intestinal epithelial cell cultures, as well as allergic mouse enterocytes.⁷⁷ Overall, enhanced transepithelial antigen transport is mediated by IgE/CD23 and regulated by IL-4 (Figure 1).

Recent studies indicated that various isoforms of CD23 mediated bidirectional transport of IgE across the epithelium in allergic murine and human intestines. DNA sequencing revealed the presence of classical and alternative CD23b transcripts lacking

exon 5 (b Δ 5) or 6 (b Δ 6) in mouse enterocytes, all of which were translated into functional IgE receptors with distinct endocytic properties.^{78,83} Mouse intestinal epithelial CD23b Δ 5 transcripts mediated apical to basolateral transport of free IgE, whereas classical CD23b transcripts displayed higher efficiency in the transcytosis of IgE/allergen complexes.^{78,83} Studies using primary human intestinal epithelial cells and transformed cell lines have also shown that CD23 transports IgE in both the mucosal-to-serosal and serosal-to-mucosal directions.⁸⁴ Both CD23 isoforms *a* and *b* transfected into human intestinal epithelial cells transcytosed bidirectionally by IgE. However, CD23a transport IgE/antigen complex was faster than CD23b in the apical-to-basolateral direction.⁸⁵ Despite the controversy over which isoform is expressed in human enterocytes,^{84,85} IgE/CD23 is responsible for enhanced transepithelial antigen uptake that results in mast cell activation. It is noteworthy that food allergens transported *via* the intestinal epithelial IgE/CD23 are protected from lysosomal degradation, and therefore, intact antigenic forms of the proteins are preserved.⁸⁶ Taken together, these findings suggest that epithelial CD23 is involved in the bidirectional transport of IgE that accounts for the high luminal concentration of IgE in allergic situations and the enhanced uptake of IgE-bound antigen complexes leading to anaphylactic responses.

6. Novel Therapeutic Interventions

Currently dietary exclusion is still the most effective measure for the prevention of allergic sensitization and anaphylactic responses in high-risk children and adults. Novel immunotherapy by delivering allergen via subcutaneous or sublingual routes have shown some degree of symptom alleviation in experimental models and human studies, yet large scale blinded, placebo-controlled trials are needed to confirm the effects for long-term desensitization.⁸⁷ The use of anti-IgE antibody increased the threshold of sensitivity to allergens in a double-blind, randomized, dose-ranging trials for peanut allergic patients.⁸⁸ Moreover, targeting CD23 with monoclonal antibody has been shown to decrease total serum IgE levels in approximately 75% of patients in phase I clinical trial of allergic asthma and was proposed as candidate therapy for treating allergic diseases for subgroups of patients.⁸⁹ Additional information is needed to develop safe and effective treatments for food allergy. The understanding of the molecular mechanism underlying epithelial barrier defects and induction of allergy may shed light to the advance in clinical management of atopic disorders.

References

- Ross MP, Ferguson M, Street D, Klontz K, Schroeder T, Luccioli S. Analysis of food-allergic and anaphylactic events in the National Electronic Injury Surveillance System. *J Allergy Clin Immunol* 2008;121:166–71.
- De S, I, Mehr SS, Tey D, Tang ML. Paediatric anaphylaxis: a 5 year retrospective review. *Allergy* 2008;63:1071–6.
- Herz U. Immunological basis and management of food allergy. *J Pediatr Gastroenterol Nutr* 2008;47(Suppl 2):54–7.
- Husby S. Food allergy as seen by a paediatric gastroenterologist. *J Pediatr Gastroenterol Nutr* 2008;47(Suppl 2):49–52.
- Wills-Karp M, Santeliz J, Karp CL. The germless theory of allergic disease: revisiting the hygiene hypothesis. *Nature Rev Immunol* 2001;1:69–75.
- Strachan DP. Hay fever, hygiene, and household size. *BMJ* 1989;299:1259–60.
- Bloomfield SF, Stanwell-Smith R, Crevel RW, Pickup J. Too clean, or not too clean: the hygiene hypothesis and home hygiene. *Clin Exp Allergy* 2006;36:402–25.
- Faria AM, Weiner HL. Oral tolerance. *Immunol Rev* 2005;206:232–59.
- Asai K, Hachimura S, Kimura M, et al. T cell hyporesponsiveness induced by oral administration of ovalbumin is associated with impaired NFAT nuclear translocation and p27kip1 degradation. *J Immunol* 2002;169:4723–31.
- Keita AV, Gullberg E, Ericson AC, et al. Characterization of antigen and bacterial transport in the follicle-associated epithelium of human ileum. *Lab Invest* 2006;86:504–16.
- Chabot S, Wagner JS, Farrant S, Neutra MR. TLRs regulate the gatekeeping functions of the intestinal follicle-associated epithelium. *J Immunol* 2006;176:4275–83.
- Fujihashi K, Dohi T, Rennert PD, et al. Peyer's patches are required for oral tolerance to proteins. *Proc Natl Acad Sci USA* 2001;98:3310–5.
- Spahn TW, Fontana A, Faria AM, et al. Induction of oral tolerance to cellular immune responses in the absence of Peyer's patches. *Eur J Immunol* 2001;31:1278–87.
- Spahn TW, Weiner HL, Rennert PD, et al. Mesenteric lymph nodes are critical for the induction of high-dose oral tolerance in the absence of Peyer's patches. *Eur J Immunol* 2002;32:1109–13.
- Spahn TW, Kucharzik T. Modulating the intestinal immune system: the role of lymphotoxin and GALT organs. *Gut* 2004;53:456–65.
- Tsuji NM, Mizumachi K, Kurisaki J. Antigen-specific, CD4+CD25+regulatory T cell clones induced in Peyer's patches. *Int Immunol* 2003;15:525–34.
- Ishikawa H, Tanaka K, Maeda Y, et al. Effect of intestinal microbiota on the induction of regulatory CD25+CD4+T cells. *Clin Exp Immunol* 2008;153:127–35.
- Frossard CP, Tropia L, Hauser C, Eigenmann PA. Lymphocytes in Peyer patches regulate clinical tolerance in a murine model of food allergy. *J Allergy Clin Immunol* 2004;113:958–64.
- Tsuji NM, Mizumachi K, Kurisaki J. Interleukin-10-secreting Peyer's patch cells are responsible for active suppression in low-dose oral tolerance. *Immunology* 2001;103:458–64.
- Miller A, Lider O, Roberts AB, Sporn MB, Weiner HL. Suppressor T cells generated by oral tolerization to myelin basic protein suppress both in vitro and in vivo immune responses by the release of transforming growth factor beta after antigen-specific triggering. *Proc Natl Acad Sci USA* 1992;89:421–5.
- Chung Y, Lee SH, Kim DH, Kang CY. Complementary role of CD4+CD25+regulatory T cells and TGF-beta in oral tolerance. *J Leukoc Biol* 2005;77:906–13.
- Ando T, Hatsushika K, Wako M, et al. Orally administered TGF-beta is biologically active in the intestinal mucosa and enhances oral tolerance. *J Allergy Clin Immunol* 2007;120:916–23.
- Zhang X, Izikson L, Liu L, Weiner HL. Activation of CD25(+) CD4(+) regulatory T cells by oral antigen administration. *J Immunol* 2001;167:4245–53.
- Sun JB, Raghavan S, Sjolting A, Lundin S, Holmgren J. Oral tolerance induction with antigen conjugated to cholera toxin B subunit generates both Foxp3+CD25+and Foxp3-. *J Immunol* 2006;177:7634–44.
- Coombes JL, Siddiqui KR, Rancibia-Carcamo CV, et al. A functionally specialized population of mucosal CD103+DCs induces Foxp3+regulatory T cells via a TGF-beta and retinoic acid-dependent mechanism. *J Exp Med* 2007;204:1757–64.
- Roth-Walter F, Berin MC, Arnaboldi P, et al. Pasteurization of milk proteins promotes allergic sensitization by enhancing uptake through Peyer's patches. *Allergy* 2008;63:882–90.
- Man AL, Bertelli E, Regoli M, Chambers SJ, Nicoletti C. Antigen-specific T cell-mediated apoptosis of dendritic cells is impaired in a mouse model of food allergy. *J Allergy Clin Immunol* 2004;113:965–72.
- O'Hara AM, Shanahan F. The gut flora as a forgotten organ. *EMBO Rep* 2006;7:688–93.
- Cani PD, Delzenne NM. Gut microflora as a target for energy and metabolic homeostasis. *Curr Opin Clin Nutr Metab Care* 2007;10:729–34.
- Kelly D, King T, Aminov R. Importance of microbial colonization of the gut in early life to the development of immunity. *Mutat Res* 2007;622:58–69.
- Kelly D, Conway S. Bacterial modulation of mucosal innate immunity. *Mol Immunol* 2005;42:895–901.
- Moreau MC, Ducluzeau R, Guy-Grand D, Muller MC. Increase in the population of duodenal immunoglobulin A plasmacytes in axenic mice associated with different living or dead bacterial strains of intestinal origin. *Infect Immun* 1978;21:532–9.
- Borsutzky S, Cazac BB, Roes J, Guzmán CA. TGF-beta receptor signaling is critical for mucosal IgA responses. *J Immunol* 2004;173:3305–7.
- Mora JR, Iwata M, Eksteen B, et al. Generation of gut-homing IgA-secreting B cells by intestinal dendritic cells. *Science* 2006;314:1157–60.
- Johansen FE, Pekna M, Norderhaug IN, et al. Absence of epithelial immunoglobulin A transport, with increased mucosal leakiness, in polymeric immunoglobulin receptor/secretory component-deficient mice. *J Exp Med* 1999;190:915–22.
- Van der Waaij LA, Limburg PC, Mesander G, Van der Waaij D. In vivo IgA coating of anaerobic bacteria in human feces. *Gut* 1996;38:348–54.
- Rey J, Garin N, Spertini F, Corthésy B. Targeting of secretory IgA to Peyer's patch dendritic and T cells after transport by intestinal M cells. *J Immunol* 2004;172:3026–33.
- Kadaoui KA, Corthésy B. Secretory IgA mediates bacterial translocation to dendritic cells in mouse Peyer's patches with restriction to mucosal compartment. *J Immunol* 2007;179:7751–7.
- Massacand JC, Kaiser P, Ernst B, et al. Intestinal bacteria condition dendritic cells to promote IgA production. *PLoS One* 2008;3:2588.
- Macpherson AJ, Slack E. The functional interactions of commensal bacteria with intestinal secretory IgA. *Curr Opin Gastroenterol* 2007;23:673–8.
- Corthésy B. Roundtrip ticket for secretory IgA: role in mucosal homeostasis? *J Immunol* 2007;178:17–32.
- Sudo N, Sawamura S, Tanaka K, Aiba Y, Kubo C, Koga Y. The requirement of intestinal bacterial flora for the development of an IgE production system fully susceptible to oral tolerance induction. *J Immunol* 1997;159:1739–45.
- Tanaka K, Ishikawa H. Role of intestinal bacterial flora in oral tolerance induction. *Histol Histopathol* 2004;19:907–14.

44. Oyama N, Sudo N, Sogawa H, Kubo C. Antibiotic use during infancy promotes a shift in the T(H)1/T(H)2 balance toward T(H)2-dominant immunity in mice. *J Allergy Clin Immunol* 2001;107:153–9.
45. Maeda Y, Noda S, Tanaka K, et al. The failure of oral tolerance induction is functionally coupled to the absence of T cells in Peyer's patches under germfree conditions. *Immunobiology* 2001;204:442–57.
46. Yang PC, Jury J, Soderholm JD, Sherman PM, McKay DM, Perdue MH. Chronic psychological stress in rats induces intestinal sensitization to luminal antigens. *Am J Pathol* 2006;168:104–14.
47. Campbell DE, Hill DJ, Kemp AS. Enhanced IL-4 but normal interferon-gamma production in children with isolated IgE mediated food hypersensitivity. *Pediatr Allergy Immunol* 1998;9:68–72.
48. Hauer AC, Breese EJ, Walker-Smith JA, MacDonald TT. The frequency of cells secreting interferon-gamma and interleukin-4, -5, and -10 in the blood and duodenal mucosal of children with cow's milk hypersensitivity. *Pediatr Res* 1997;42:629–38.
49. Paul WE. Interleukin 4: a prototypic immunoregulatory lymphokine. *J Am Soc Hematol* 1991;77:1859–70.
50. Belut D, Moneret-Vautrin DA, Nicholas JP, Grilliat JP. IgE levels in intestinal juice. *Dig Dis Sci* 1980;25:323–32.
51. Brown WR, Lee EH. Studies on IgE in human intestinal fluids. *Int Arch Allergy Appl Immunol* 1976;50:87–94.
52. Negrao-Correa D, Adams LS, Bell RG. Intestinal transport and catabolism of IgE. *J Immunol* 1996;157:4037–44.
53. Ramaswamy K, Hakimi J, Bell RG. Evidence for an interleukin 4-inducible immunoglobulin E uptake and transport mechanism in the intestine. *J Exp Med* 1994;180:1793–803.
54. Yu LCH, Perdue MH. Role of mast cells in intestinal mucosal function: studies in models of hypersensitivity and stress. *Immunol Rev* 2001;179:61–73.
55. Perdue MH, Masson S, Wershil BK, Galli SJ. Role of mast cells in ion transport abnormalities associated with intestinal anaphylaxis. *J Clin Invest* 1991;87:687–93.
56. Yu LCH, Perdue MH. Immunologically mediated transport of ions and macromolecules. *Ann New York Acad Sci* 2001;915:247–59.
57. Crowe SE, Sestini P, Perdue MH. Allergic reactions of rat jejunal mucosa: Ion transport responses to luminal antigen and inflammatory mediators. *Gastroenterology* 1990;99:74–82.
58. Castro GA, Harari Y, Russel D. Mediators of anaphylaxis-induced ion transport changes in small intestine. *Am J Physiol* 1987;253:540–8.
59. Wasserman SI, Barrett KE, Huott PA, Beuerlein G, Kagnoff MF, Dharmasathaphorn K. Immune-related intestinal Cl⁻ secretion. I. Effect of histamine on the T84 cell line. *Am J Physiol* 1988;254:53–62.
60. Hardcastle J, Hardcastle PT. The secretory actions of histamine in rat small intestine. *J Physiol* 1987;388:521–32.
61. Kimberg DV, Field M, Gershon E, Henderson A. Effects of prostaglandins and cholera enterotoxin on intestinal mucosal cyclic AMP accumulation. Evidence against an essential role for prostaglandins in the action of toxin. *J Clin Invest* 1974;53:941–9.
62. Schmitz H, Fromm M, Bode H, Scholz P, Riecken EO, Schulzke JD. Tumor necrosis factor- α induces Cl⁻ and K⁺ secretion in human distal colon driven by prostaglandin E₂. *Am J Physiol* 1996;271:669–74.
63. Beubler E, Bukhave K, Rask-Madsen J. Significance of calcium for the prostaglandin E₂-mediated secretory response to 5-hydroxytryptamine in the small intestine of the rat in vivo. *Gastroenterology* 1986;90:1972–7.
64. Heyman M, Ducroc R, Desjeux JF, Morgat JL. Horseradish peroxidase transport across adult rabbit jejunum in vitro. *Am J Physiol* 1982;242:558–64.
65. Fujita M, Reinhart F, Neutra M. Convergence of apical and basolateral endocytic pathways at apical late endosomes in absorptive cells of suckling rat ileum in vivo. *J Cell Sci* 1990;97:385–94.
66. Laiping SA, Pelton-Henrion K, Small G, et al. Antigen uptake and trafficking in human intestinal epithelial cells. *Dig Dis Sci* 2000;45:1451–61.
67. Clark JA, Coopersmith CM. Intestinal crosstalk: a new paradigm for understanding the gut as the "motor" of critical illness. *Shock* 2007;28:384–93.
68. Teshima CW, Meddings JB. The measurement and clinical significance of intestinal permeability. *Curr Gastroenterol Rep* 2008;10:443–9.
69. Bojarski C, Weiske J, Schoneberg T, et al. The specific fates of tight junction proteins in apoptotic epithelial cells. *J Cell Sci* 2004;117:2097–107.
70. Utech M, Ivanov AI, Samarina SN, et al. Mechanism of IFN-gamma-induced endocytosis of tight junction proteins: myosin II-dependent vacuolarization of the apical plasma membrane. *Mol Biol Cell* 2005;16:5040–52.
71. Yu LCH, Flynn AN, Turner JR, Buret AG. SGLT-1-mediated glucose uptake protects intestinal epithelial cells against LPS-induced apoptosis and barrier defects: a novel cellular rescue mechanism? *FASEB J* 2005;19:1822–35.
72. Shen L, Black ED, Witkowski ED, et al. Myosin light chain phosphorylation regulates barrier function by remodeling tight junction structure. *J Cell Sci* 2006;119:2095–106.
73. Hsiao JK, Huang C-Y, Lu Y-Z, Yang C-Y, Yu LCH. Magnetic resonance imaging detects intestinal barrier dysfunction in a rat model of acute mesenteric ischemia/reperfusion injury. *Invest Radiol* 2009;44:329–35.
74. Berin MC, Kiliaan AJ, Yang PC, Groot JA, Taminiau JAJM, Perdue MH. Rapid transepithelial antigen transport in rat jejunum: impact of sensitization and the hypersensitivity reaction. *Gastroenterology* 1997;113:1–9.
75. Yang PC, Berin MC, Yu LCH, Conrad DH, Perdue MH. Enhanced intestinal transepithelial transport in allergic rats is mediated by IgE and CD23 (Fc ϵ RII). *J Clin Invest* 2000;106:879–86.
76. Berin MC, Kiliaan AJ, Yang PC, Groot JA, Kitamura Y, Perdue MH. The influence of mast cells on pathways of transepithelial antigen transport in rat intestine. *J Immunol* 1998;161:2561–6.
77. Yu LCH, Yang PC, Berin MC, et al. Enhanced transepithelial antigen transport in intestine of allergic mice is mediated by IgE/CD23 and regulated by interleukin-4. *Gastroenterology* 2001;121:370–81.
78. Yu LCH, Montagnac G, Yang PC, Conrad DH, Benmerah A, Perdue MH. Intestinal epithelial CD23 mediates enhanced antigen transport allergy: evidence for novel splice forms. *Am J Physiol Gastrointest Liver Physiol* 2003;285:223–34.
79. Bonnefoy JY, Lecoanet-Henchoz S, Aubry JP, Gauchat JF, Graber P. CD23 and B-cell activation. *Curr Biol* 1995;7:355–9.
80. Ford JW, Kilmon MA, Haas KM, Shelburne AE, Chan-Li Y, Conrad DH. In vivo murine CD23 destabilization enhances CD23 shedding and IgE synthesis. *Cell Immunol* 2006;243:107–17.
81. Zhang M, Murphy RF, Agrawal DK. Decoding IgE Fc receptors. *Immunol Res* 2007;37:1–16.
82. Kaiserlian D, Lachaux A, Gosjean I, Graber P, Bonnefoy JY. Intestinal epithelial cells express the CD23/Fc ϵ RII molecule: enhances expression in enteropathies. *Immunology* 1993;80:90–5.
83. Montagnac G, Yu LC, Bevilacqua C, et al. Differential role for CD23 splice forms in apical to basolateral transcytosis of IgE/allergen complexes. *Traffic* 2005;6:230–42.

84. Tu Y, Salim S, Bourgeois J, et al. CD23-mediated IgE transport across human intestinal epithelium: inhibition by blocking sites of translation or binding. *Gastroenterology* 2005;129:928–40.
85. Li H, Nowak-Węgrzyn A, Charlop-Powers Z, et al. Transcytosis of IgE-antigen complexes by CD23a in human intestinal epithelial cells and its role in food allergy. *Gastroenterology* 2006;131:47–58.
86. Bevilacqua C, Montagnac G, Benmerah A, et al. Food allergens are protected from degradation during CD23-mediated transepithelial transport. *Int Arch Allergy Immunol* 2004;135:108–16.
87. Skripak JM, Sampson HA. Towards a cure for food allergy. *Curr Opin Immunol* 2008;20:690–6.
88. Leung DY, Sampson HA, Yunginger JW, et al. Effect of anti-IgE therapy in patients with peanut allergy. *N Engl J Med* 2003;348:986–93.
89. Rosenwasser LJ, Meng J. Anti-CD23. *Clin Rev Allergy Immunol* 2005;29:61–72.